Severity Measure for Social Anxiety Disorder (Social Phobia)—Adult

Age: ____ Sex: Male ☐ Female ☐ Date:

socio part	ructions: The following questions ask about al situations. Usual social situations include ies, introducing yourself to others, having thers, and eating and writing in public. Ple	e: public conversa	speaking, speations, giving ar	aking in mee nd receiving	tings, atten complimen	ding social ts, making i	events or requests
34		ciin					Clinician Use
	During the PAST 7 DAYS, I have	Never	Occasionally	Half of the time	Most of the time	All of the time	Item score
1.	felt moments of sudden terror, fear, or fright in social situations	0 0	0 1	□ 2	□ 3	4	la constant
2.	felt anxious, worried, or nervous about social situations	0 0	1	□ 2	□ 3	4	
3.	had thoughts of being rejected, humiliated, embarrassed, ridiculed, or offending others	00	□ 1	□ 2	3	4	
4.	felt a racing heart, sweaty, trouble breathing, faint, or shaky in social situations	0	01	2	3	4	
5.	felt tense muscles, felt on edge or restless, or had trouble relaxing in social situations	00	0 1	a 2	3	4	5 6734
6.	avoided, or did not approach or enter, social situations	0 0	0 1	Q 2	□ 3	4	Water.
7.	left social situations early or participated only minimally (e.g., said little, avoided eye contact)	0 0	0 1	a 2	3	4	
8.	spent a lot of time preparing what to say or how to act in social situations	00	0 1	a 2	3	4	-
9.	distracted myself to avoid thinking about social situations	0 0	0 1	□ 2	3	4	
10.	needed help to cope with social situations (e.g., alcohol or medications, superstitious objects)	0	0 1	a 2	3	4	
	and the same of the same of the same of				Total/Partial		
Prorated Total Raw Score: (if 1-2 items left unanswered) Average Total Score:							

Craske M, Wittchen U, Bogels S, Stein M, Andrews G, Lebeu R. Copyright © 2013 American Psychiatric Association. All rights reserved.

This material can be reproduced without permission by researchers and by clinicians for use with their patients.